# **HPLC Application**

ID No.: 19680



Kinetex<sup>®</sup>

## Estrone, Estradiol, and Estriol from Water @ pH 2.0 Strata-X via Kinetex XB-C18 150x4.60mm

Kinetex® 2.6  $\mu$ m XB-C18 100 Å, LC Column 150 x 4.6 mm, Ea

150 x 4.6 mm ID **Dimensions:** Order No: 00F-4496-E0 **Elution Type:** Gradient

6.9mM Ammonium Acetate pH 4.0 with 25% Acetonitrile (v/v) Eluent A:


Eluent B:

Gradient Step No. Time (min) Pct A Pct B **Profile:** 0 100 0 2 10 35 65

Flow Rate: 1.2 mL/min Col. Temp.: ambient

**Detection:** UV-Vis Abs.-Variable Wave.(UV) @ 230 nm (ambient)





### **ANALYTES:**

- Estriol 1
- 2 Ethyl paraben
- 17-beta-Estradiol
- Estrone



# **Sample Preparation Details**

for **HPLC** Application ID No.: **19680** 



## Estrone, Estradiol, and Estriol from Water @ pH 2.0 Strata-X via Kinetex XB-C18 150x4.60mm

### **PRODUCT DESCRIPTION:**

Strata™-X 33 µm Polymeric Reversed Phase, 200 mg / 6 mL, Tubes , 30/Pk

Order No.: 8B-S100-FCH

### **SOLID PHASE EXTRACTION (SPE) PRODCEDURE:**

**Note:** The solvent volumes shown below are for a 200 mg bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:
Load:
Wash:
Dry:
1 Minute
Elute:
Final Prep and Analysis:
Inject, Oul on HDICIIVIVIC Abo Variable Ways (IIV) @ 220 nm (ambient)

Inject: 0 µL on HPLC UV-Vis Abs.-Variable Wave.(UV) @ 230 nm (ambient)

ANALYTES:		Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1	Estriol	30			81	
2	Ethyl paraben	20			100	
3	17-beta-Estradiol	30			98	
4	Estrone	30			88	

This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com support@phenomenex.com