HPLC Application

ID No.: 20926



JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B

Kinetex® 2.6 µm C18 100 Å, LC Column 150 x 3 mm, Ea

150 x 3 mm ID **Dimensions:** Order No: 00F-4462-Y0 **Elution Type:** Gradient

Eluent A: 10mM ammonium formate

Eluent B: Acetonitrile

Gradient	Step No.	Time (min)	Pct A	Pct B
Profile:	1	0	55	45
	2	7	50	50
	3	7.01	5	95
	4	10	5	95



Products used in this application:



Flow Rate: 0.6 mL/min Col. Temp.: ambient

Mass Spectrometer (MS) @ amu (ambient) **Detection:**

Detector Info: <a target="_blank"

href="https://sciex.com/products/mass-spectrometers?utm_campaign=2019%20application%20search&utm_source=phenomenex&utm_medium=referral">SCIEXPositive


7.00

CAD: CUR: GS1:

Analyst Note:

20.00 50.00 GS2: 50.00 5500.00 IS: 550.00 TEM: ihe: ON ΕP 10.00

App ID 20926



©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



HPLC Application ID No.: **20926**



JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B **ANALYTES:**

- JWH-073-Butanoic acid metabolite
- JWH-018-5-Hydroxypentyl metabolite
- JWH-018-4-Hydroxypentyl metabolite



Sample Preparation Details

for HPLC Application ID No.: 20926



JHW018-hydroxypentyl isomers in Urine using Strata-X-Drug B

PRODUCT DESCRIPTION:

Strata™-X-Drug B 33 µm Polymeric Strong Cation, 60 mg / 3 mL, Tubes , 50/Pk

Order No.: 8B-S128-UBJ

SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

Note: The solvent volumes shown below are for a 60 mg bed mass.
The solvent volumes will need to be adjusted for a smaller or larger bed mass.
Condition:
Load:
Sample Hydrolysis: Combine 1 mL Human Urine sample (spiked with analytes at 50 ng/mL), 2 mL of 100 mM sodium acetate buffer, pH 5.0, 25 uL -D-Glucoronidase (Patella Vulgata from Sigma, 100KU).
Vortex 10-15 secs, followed by incubation for 2 hours in a shaker at 55o C to complete hydrolysis of the glucuronides.
Wash:
Dry:
15 mins under 10-15" of Hg
Elute:
Final Prep and Analysis:
Extraction Protocol (No need for conditioning or equilibration of the cartridge)
Inject: 10 µL on HPLC Mass Spectrometer (MS) @ amu (ambient)

ANALYTES: Spiked Conc. % Rec %RSC Log P pKa (ng/mL) (n=0)

- 1 JWH-073-Butanoic acid metabolite 0
- JWH-018-5-Hydroxypentyl metabolote
- **3** JWH-018-4-Hydroxypentyl metabol**0**te

This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com

