HPLC Application

ID No.: **22785**



Automated Extraction of Prednisolone & Cortisone from Plasma Using Novum SLE & Kinetex 2.6u Biphenyl

Kinetex® 2.6 μm Biphenyl 100 Å, LC Column 50 x 3.0 mm, Ea

50 x 3 mm ID **Dimensions:** Order No: 00B-4622-Y0 Elution Type: Gradient

Eluent A: 10 mm ammonium acetate in water Eluent B: 10 mm ammonium acetate in methanol

Gradient	Step No.	Time (min)	Pct A	Pct B
Profile:	1	0	50	50
	2	2	5	95
	3	3.1	5	95
	4	3.11	50	50
	5	5	50	50



Products used in this application:




Flow Rate: 0.45 mL/min

Col. Temp.: 25 °C

Detection: Tandem Mass Spec (MS-MS) @ (ambient)

Detector Info:



ANALYTES:

Cortisone

Retention Time: 2.66 min

Cortisol-D4 (Internal standard)

Retention Time: 2.69 min

Prednisolone 3

Retention Time: 2.92 min

@2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Sample Preparation Details for HPLC Application ID No.: 22785



Automated Extraction of Prednisolone & Cortisone from Plasma Using Novum SLE & Kinetex 2.6u Biphenyl

PRODUCT DESCRIPTION:

Novum SLE MINI 96-Well Plate, 1/Pk

Order No.: 8E-S138-FGA

SOLID PHASE EXTRACTION (SPE) PRODCEDURE:

Note: The solvent volumes shown below are for a 200 mg bed mass.
The solvent volumes will need to be adjusted for a smaller or larger bed mass.
Condition:
Load:
Sample pre-treatment:
Dilute 150 uL of human plasma (spiked with 25 ng/mL and 125 ng/mL of cortisone and prednisolone respectively) with 150 uL of 50 mM sodium phosphate dibasic heptahydrate, pH unadjusted.
Mix briefly (3-5 sec).
Sample loading: Load the sample from pre-treatment step above onto the Novum plate and apply a short and gentle pulse of vacuum (~ 10" of Hq for 20 secs) until the sample has completely entered the media.
Load the sample from pre-deathert step above onto the novum piace and apply a short and gende pulse of vacuum (** 10* of rig to 20 secs) that the sample has completely entered the media.
Wait for 5 minutes
Wash:
Dry:
Elute:
Final Prep and Analysis:
Sample pre-treatment
Dilute 150 uL of human plasma (spiked with 25 ng/mL and 125 ng/mL of cortisone and
Inject: 10 μL on HPLC Tandem Mass Spec (MS-MS) @ (ambient)

ANAL	LYTES:	(ng/mL)	Log P	рка	% Rec	%RSC (n=0)
1	Cortisone	25			99	
2	Cortisol-D4 (Internal standard)	50				
3	Prednisolone	125			95	

This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals. Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com

