

Chiral Separation of Dextromethorphan/Levomethorphan and Dextrorphan/Levorphanol on Lux 3u AMP, 150x4.6



Column: Lux® 3 µm AMP, LC Column 150 x 4.6 mm, Ea
Dimensions: 150 x 4.6 mm ID
Order No: 00F-4751-E0
Elution Type: Gradient
Eluent A: 5mM ammonium bicarbonate, adjusted to pH 11 with ammonium hydroxide
Eluent B: Acetonitrile

Gradient Profile:	Step No.	Time (min)	Pct A	Pct B
	1	0	55	45
	2	4.5	55	45
	3	4.6	20	80
	4	8	20	80
	5	8.1	55	45
	6	10	55	45

Flow Rate: 1 mL/min
Col. Temp.: 40 °C
Detection: LC/MS/MS @ (ambient)
Analyst Note: Urine hydrolyzed with beta-glucuronidase and extracted with Strata-X-C



Products used in this application:



ANALYTES:

- Dextrorphan 1
Retention Time: 3.85 min
- Dextrorphan 2
Retention Time: 3.85 min
- Levorphanol 1
Retention Time: 4.17 min
- Levorphanol 2
Retention Time: 4.17 min
- Dextromethorphan 1
Retention Time: 7.24 min
- Dextromethorphan 2
Retention Time: 7.24 min
- Levomethorphan 1
Retention Time: 7.69 min
- Levomethorphan 2
Retention Time: 7.69 min

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com

support@phenomenex.com

Sample Preparation Details

for HPLC Application ID No.: 25783

Chiral Separation of Dextromethorphan/Levomethorphan and Dextrorphan/Levorphanol on Lux 3u AMP, 150x4.6

PRODUCT DESCRIPTION:

Strata™-X-C 33 µm Polymeric Strong Cation, 30 mg / well, 96-Well Plates , 2/Pk

Order No.: 8E-S029-TGB

SOLID PHASE EXTRACTION (SPE) PROCEDURE:

Note: The solvent volumes shown below are for a bed mass.

The solvent volumes will need to be adjusted for a smaller or larger bed mass.

Condition:

Load:

Wash:

Dry:

Elute:

Final Prep and Analysis:

Urine sample pretreatment: Combine 200µL urine sample, 200µL 100mM ammonium acetate buffer (pH 4.0), 20µL internal standard solution (250ng/mL), and 20µL of beta-glucuronidase

Inject: 10 µL on HPLC LC/MS/MS @ (ambient)

ANALYTES:	Spiked Conc. (ng/mL)	Log P	pKa	% Rec	%RSC (n=0)
1 Dextrorphan 1	0				
2 Dextrorphan 2	0				
3 Levorphanol 1	0				
4 Levorphanol 2	0				
5 Dextromethorphan 1	0				
6 Dextromethorphan 2	0				
7 Levomethorphan 1	0				
8 Levomethorphan 2	0				

Note: This method is designed as a convenient starting point for further investigation and can be tailored to meet your extraction goals.
Call your local Phenomenex Representative for assistance in method development and optimization techniques.

©2025 Phenomenex Inc. All rights reserved.

For more information contact your Phenomenex Representative at support@phenomenex.com



Phenomenex products are available worldwide.

www.phenomenex.com

support@phenomenex.com